Recombinant Alpha-2-Heremans Schmid Glycoprotein (AHSG) Instruction Manual

SIPA332Po01

Sus scrofa; Porcine (Pig)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass40.4kDa

Accurate Molecular Mass 50kDa(Analysis of differences refer to the manual)

Residues & Tags Val16~Ile362 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 5.5

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

VPHGP ILGYREPACD DVETEQAALA AVDYINKHLP
RGYKHTLNQV DSVKVWPRRP AGEVFDIEID TLETTCHVLD PTPLANCSVR
QLTEHAVEGD CDFHVLKQDG QFSVLFAKCD SSPDSAEDVH KVCPNCPLLA
PLNDSRVVHA AESALAAFNA QSNGSYLQLV EISRAQLVPL SASVSVEFAV
AVTDCVAKEA YSPTKCNLLV EKQYGFCKGT VTAKVNEEDV AVTCTVFQTQ
PVVLQPQPAG ADAGATPVVD AAATASPLAD VPAASLVVGP MVVAVPPGIP
PVHRSHYDLR HSFSGVASVE SASGEAFHVG KTPKGAQPSI PAADGSVPVV
RPCPGRIRHF KI

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.