

# Recombinant Coagulation Factor V (F5) Instruction Manual

**SIPA352Hu03**

**Homo sapiens (Human)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Secreted
<b>Predicted Molecular Mass</b>	35.2kDa
<b>Accurate Molecular Mass</b>	34kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Arg1908~Val2203 with N-terminal His Tag
<b>Buffer Formulation</b>	PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 95%
<b>Isoelectric Point</b>	9.5
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## SEQUENCE

```
RMP MGLSTGIISD SQIKASEFLG YWEPRLARLN NGGSYNAWSV
EKLAAEFASK PWIQVDMQKE VIITGIQTQG AKHYLKSCYT TEFYVAYSSN
QINWQIFKGN STRNVMYFNG NSDASTIKEN QFDPPIVARY IRISPTRAYN
RPTLRLELQG CEVNGCSTPL GMENKGIENK QITASSFKKS WWDYWEPRF
ARLNAQGRVN AWQAKANNK QWLEIDLLKI KKITAIITQG CKSLSEMYV
KSYTIHYSEQ GVEWKPYRLK SSMVDKIFEG NTNTKGHVKN FFPPIISRF
IRV
```

## USAGE

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.


## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **Image**



SDS-PAGE Image

## **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.