

Recombinant Homing Associated Cell Adhesion Molecule (HCAM) Instruction Manual

SIPA451Mu03

Mus musculus (Mouse)

| | |
|---------------------------------|---|
| Source | Prokaryotic expression |
| Host | E.coli |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | Membrane |
| Predicted Molecular Mass | 56.2kDa |
| Accurate Molecular Mass | 60kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Gln23~Pro258 with N-terminal His and GST Tag |
| Buffer Formulation | PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. |
| Traits | Freeze-dried powder |
| Purity | > 80% |
| Isoelectric Point | 5.1 |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

SEQUENCE

```
QIDLNVTC RYAGVFHVEK NGRYSISRTE  
AADLCQAFNS TLPTMDQMKL ALSKGFETCR YGFIEGNVVI PRIHPNAICA  
ANHTGVYILV TSNTSHYDTY CFNASAPPEE DCTSVTDLPN SFDGPVTITI  
VNRDGTRYSK KGEYRTHQED IDASNIIDDD VSSGSTIEKS TPEGYILHTY  
LPTEQPTGDQ DDSFFIRSTL ATIASTVHVK SHAAAQKQNN WIWSWFGNSQ  
STTQTQEP
```

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.