# Recombinant Heat Shock 70kDa Protein 1B (HSPA1B) Instruction Manual

# SIPA510Bo01

## Bos taurus; Bovine (Cattle)

Source	Prokaryotic expression		
Host	E.coli		
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)		
Subcellular Location	Secreted		
Predicted Molecular Mass	71.5kDa		
Accurate Molecular Mass	70kDa(Analysis of differences refer to the manual)		
Residues & Tags	Met1~Asp641 with N-terminal His Tag		
<b>Buffer Formulation</b>	PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.		
Traits	Freeze-dried powder		
Purity	> 95%		
Isoelectric Point	5.9		
Applications	Positive Control; Immunogen; SDS-PAGE; WB.		

# SEQUENCE

MAKNTAIGID	LGTTYSCVGV	FQHGKVEIIA	NDQGNRTTPS	YVAFTDTERL
IGDAAKNQVA	LNPQNTVFDA	KRLIGRKFGD	PVVQSDMKHW	PFRVINDGDK
PKVQVSYKGE	TKAFYPEEIS	SMVLTKMKEI	AEAYLGHPVT	NAVITVPAYE
NDSQRQATKD	AGVIAGLNVL	RIINEPTAAA	IAYGLDRTGK	GERNVLIFDL
GGGTFDVSIL	TIDDGIFEVK	ATAGDTHLGG	EDFDNRLVNH	FVEEFKRKHK
KDISQNKRAV	RRLRTACERA	KRTLSSSTQA	SLEIDSLFEG	IDFYTSITRA
RFEELCSDLF	RSTLEPVEKA	LRDAKLDKAQ	IHDLVLVGGS	TRIPKVQKLL
QDFFNGRDLN	KSINPDEAVA	YGAAVQAAIL	MGDKSENVQD	LLLLDVAPLS
LGLETAGGVM	TALIKRNSTI	PTKQTQIFTT	YSDNQPGVLI	QVYEGERAMT
RDNNLLGRFE	LSGIPPAPRG	VPQIEVTFDI	DANGILNVTA	TDKSTGKANK
ITITNDKGRL	SKEEIERMVQ	EAEKYKAEDE	VQRERVSAKN	ALESYAFNMK
SAVEDEGLKG	KISEADKKKV	LDKCQEVISW	LDANTLAEKD	EFEHKRKELE
QVCNPIISRL	YQGAGGPGAG	GFGAQGPKGG	SGSGPTIEEV	D

#### USAGE


Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

#### STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

### STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.