Recombinant Heme Oxygenase 1 (HO1) Instruction Manual

SIPA520Bo01

Bos taurus; **Bovine** (Cattle)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Membrane **Predicted Molecular Mass** 36.6kDa

Accurate Molecular Mass 37kDa(Analysis of differences refer to the manual)

Residues & Tags Met1~Met289 with N-terminal His Tag

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, **Buffer Formulation**

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

> 90% **Purity Isoelectric Point** 7.9

Positive Control; Immunogen; SDS-PAGE; WB. **Applications**

SEQUENCE

MERPOPDSSM PODLSEALKE ATKEVHTOAE NAEFMKNFOK GELTOEGFKL VMASLYHIYV ALEEEIERNK ENPVYTPLYF PEELHRRASL EQDMAFWYGP RWQEAIPYTQ ATKRYVQRLQ EVGRTEPELL VAHAYTRYLG DLSGGQVLKK IAQKALNLPS SGEGLAFFTF PNIASATKFK QLYRSRMNTL EMTPEVRQRV LDEAKTAFLL NIQLFEELQG LLTQKAKDHD PLQAPELHRR AGSKVQDLAP

TKASRGKPOP SVLSOAPLLR WVLTLSFLVA TVAVGLYAM

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.





SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.