Recombinant Plasminogen Activator, Urokinase (uPA) Instruction Manual

SIPA541Po01

Sus scrofa; Porcine (Pig)

| Source | Prokaryotic expression |
|--------------------------|---|
| Host | E.coli |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | Secreted |
| Predicted Molecular Mass | 20.3kDa |
| Accurate Molecular Mass | 21kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Ser21~Pro168 (Accession # P04185) with |
| Buffer Formulation | PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. |
| Traits | Freeze-dried powder |
| Purity | > 95% |
| Isoelectric Point | 8.7 |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

USAGE


Reconstitute in PBS or others.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.