

Recombinant Podocalyxin (PODXL) Instruction Manual

SIPA592Mu02

Mus musculus (Mouse)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Membrane
Predicted Molecular Mass	43.7kDa
Accurate Molecular Mass	53kDa(Analysis of differences refer to the manual)
Residues & Tags	His22~Ser404 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	5.0
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
                HNGNETSTS AIKSSTVQSH QSATTSTEVT  
TGHPVASTLA STQPSNPTPF TTSTQSPSMP TSTPNPTSNO SGGNLTSSVS  
EVDKTKTSSP SSTAFTSSSG QTASSGGKSG DSFTTAPTTT LGLINVSSQP  
TDLNNTSKLL STPTTDNTTS PQQPVDSSPS TASHPVGQHT PAAVPSSSGS  
TPSTDNSTLT WKPTTHKPLG TSEATQPLTS QTPGITTLPV STLQQSMAS  
VGTTEEFTH LISNGTPVAP PGPSTPSPIW AFGNYQLNCE PPIRPDEELL  
ILNLTRASLC ERSPLDEKEK LVELLCHSVK ASFKPAEDLC TLHVAPILDN  
QAVAVKRIII ETKLSPKAVY ELLKDRWDDL TEAGVSDMKL GKEGPPEVNE  
DRFS
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.