

Recombinant Tissue Factor Pathway Inhibitor (TFPI) Instruction Manual

SIPA616Mu01

Mus musculus (Mouse)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	35.5kDa
Accurate Molecular Mass	36kDa(Analysis of differences refer to the manual)
Residues & Tags	Leu29~Asn306 with N-terminal His Tag
Buffer Formulation	PBS, pH7.4, containing 0.05% SKL, 5% Trehalose.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	8.8
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
LS EEADDTSEL GSMKPLHTFC  
AMKADDGPKK AMIRSYFFNM YTHQCEEFYI GGCEGNENRF DTLEECKKTC  
IPGYEKTAVK AASGAERPFD CFLEEDPGLC RGYMKRYLYN NQTKQCERFV  
YGGCLGNRNN FETLDECKKI CENPVHSPSP VNEVQMSDYV TDGNTVTDRS  
TVNNIVVPQS PKVPRRRDYR GRPWCLQPAD SGLCKASERR FYYNSATGKC  
HRFNVTGCGG NNNNFTTRRR CLRCKTGLI KNKSKGVVKI QRRKAPFVKV  
VYESIN
```

USAGE


Reconstitute in ddH₂O to a concentration of 0.1-0.2 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.