

Recombinant Plasminogen Activator Inhibitor 1 (PAI1) Instruction Manual

SIPA618Mu01

Mus musculus (Mouse)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	n/a
Predicted Molecular Mass	31.7kDa
Accurate Molecular Mass	28kDa(Analysis of differences refer to the manual)
Residues & Tags	Gln141~Pro389 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	6.2
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
QTMVKQVDFS  
EVERARFIIN DWVERHTKGM INDLLAKGAV DELTRLVLVN ALYFSGQWKT  
PFLEASTHOR LFHKSDGSTV SVPMMASNK FNYTEFTTPD GLEYDVVELP  
YQRDTLSMFI AAPFEKDVHL SALTNILDAE LIRQWGNMT RLPRLILPK  
FSLETEVDLR GPLEKLGMPD MFSATLADFT SLSDQEQLSV AQALQKVRIE  
VNESGTVASS STAFVISARM APTEMVIDRS FLFVVRHNP
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.