Recombinant Vitronectin (VTN) Instruction Manual

SIPB024Ra01

Rattus norvegicus (Rat)

Source	Prokaryotic expression		
Host	E.coli		
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)		
Subcellular Location	n/a		
Predicted Molecular Mass	47.1kDa		
Accurate Molecular Mass	55kDa(Analysis of differences refer to the manual)		
Residues & Tags	Gln21~Arg400 with N-terminal His Tag		
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.		
Traits	Freeze-dried powder		
Purity	> 90%		
Isoelectric Point	6.3		
Applications	Positive Control; Immunogen; SDS-PAGE; WB.		

SEQUENCE

		QESCKGRCTQ	GFMASKKCQC	DELCTYYQSC
CVDYMEQCKP	QVTRGDVFTM	PEDEYWSYDY	PEETKNSTNA	GVQSENTSLH
FNLKPRAEET	IKPTTPDPQE	QSNTQEPEVG	QQGVAPRPDT	TDEGTSEFPE
EELCSGKPFD	AFTDLKNGSL	FAFRGEYCYE	LDETAVRPGY	PKLIQDVWGI
EGPIDAAFTR	INCQGKTYLF	KGSQYWRFED	GVLDPDYPRN	ISEGFSGIPD
NVDAALALPA	HSYSGRERVY	FFKGKQYWEY	EFQQQPSQEE	CEGSSLSAVF
EHFALLQRDS	WENIFELLFW	GRSSDGAKGP	QFISRDWHGV	PGKVDAAMAG
RIYITGSTFR	SVQAKKQKSG	RRSRKRYRSR	RGRGHSRSRS	RSMSSRRPSR

USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.