

Recombinant Heat Shock 70kDa Protein 1A (HSPA1A) Instruction Manual

SIPB053Mu02

Mus musculus (Mouse)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Nucleus, Cytoplasm
Predicted Molecular Mass	73.8kDa
Accurate Molecular Mass	70kDa(Analysis of differences refer to the manual)
Residues & Tags	Met1~Asp641 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	5.4
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

MAKNTAIGID LCPTYSCVGV FQHGKVEIIA NDQGNRTTPS YVAFTDTERL
IGDAAKNQVA LNPQNTVFDA KRLIGRKFGD AVVQSDMKHW PFQVVNDGDK
PKVQVNQYKGE SRSFFPPEEIS SMVLTKMKEI AEAYLGHPTV NAVITVPAYF
NDSQRQATKD AGVIAGLNVL RIINEPTAAA IAYGLDRTGK GERNVLIFDL
GGGTFDVSIL TIDDGIFEVK ATAGDTHLGG EDFDNRLVSH FVEEFKRKHK
KDISQNKRRAV RRLRTACERA KRTLSSSTQA SLEIDSLFEG IDFYTSITRA
RFEELCSDLF RGTEPVEKA LRDAKMDKAQ IHDLVLVGGS TRIPKVQKLL
QDFFNGRDLN KSINPDEAVA YGAAVQAAIL MGDKSENVQD LLLL DVAPLS
LGLETAGGVM TALIKRNSTI PTKQTQTFTT YSDNQPGVLI QVYEGERAMT
RDNNILLGRFE LSGIPPAPRG VPQIEVTDFDI DANGILNVTA TDKSTGKANK
ITITNDKGRL SKEEIERMVQ EAERYKAED E VQRDRVAAKN ALESYAFNMK
SAVEDEGLKG KLSEADKKKV LDKCQEVISW LDSNTLADKE EFVHKREELE
RVCSPPIISGL YQGAGAPGAG GFQAQAPKGA SGSGPTIEEV D

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.