

# Recombinant Nucleoporin 88 (NUP88) Instruction Manual

SIPB098Ra01

*Rattus norvegicus* (Rat)

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Nucleus
<b>Predicted Molecular Mass</b>	33.2kDa
<b>Accurate Molecular Mass</b>	35kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Thr159~Ile422 with N-terminal His Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 97%
<b>Isoelectric Point</b>	5.2
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## SEQUENCE

```
TI PIAERFFTSS TSLTLKHAAW YPSEMIDPHI VLLTSDNVIR  
IYSLREPQTP TKVIVLSEAE EESLILNKGR AYTASLGETA VAFDFGPLVT  
VSKNMFQKD REAVAYPLYI LYENGETFLT YVSLHSPGN IGKLLGPLPM  
HPAAEDNYGY DACAILCLPC VPNILVIATE SGMLYHCVVL EGEEDDDQTL  
EKSWDPRADL IPSLYVFECV ELELALKLAS AEDDPFASDF SCPIKLHRDP  
KCPSRYHCSH EAGVHSVGLT WI
```

## USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.  
Do not vortex.


## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image



SDS-PAGE Image

## [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.