

Recombinant Lactate Dehydrogenase A (LDHA) Instruction Manual

SIPB118Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Cytoplasm
Predicted Molecular Mass	40.8kDa
Accurate Molecular Mass	41kDa(Analysis of differences refer to the manual)
Residues & Tags	Met1~Phe332 plus VPVC between Ser197 and Val198 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	8.1
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
MATLKDQLIY NLLKEEQTPQ NKITVVGVGA VGMACAISIL MKDLADELAL  
VDVIEDKLLG EMDLQHGSL FLRTPKIVSG KDYNVTANSK LVIITAGARQ  
QEGESRLNLV QRNVNIFKFI IPNVVKYSPN CKLLIVSNPV DILTYVAWKI  
SGFPKNRVIG SGCNLSARF RYLMGERLGV HPLSCHGWVL GEHGDSSVPV  
CVPVWSGMNV AGVSLKTLHP DLGTDKDKAQ WKEVHKQVVE SAYEVIKLLG  
YTSWAIGLSV ADLAESIMKN LRRVHPVSTM IKGLYGIKDD VFLSVPCILG  
QNGISDLVKV TLTSEEEARL KKSADTLWGI QKELQF
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.