# Recombinant Interleukin 10 Receptor Alpha (IL10Ra) Instruction Manual

## SIPB154Mu02

#### Mus musculus (Mouse)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass27.5kDa

Accurate Molecular Mass 28kDa(Analysis of differences refer to the manual)

Residues & Tags Leu17~Ile241 with N-terminal His Tag

100mMNaHCO<sub>3</sub>, 500mMNaCl, pH8.3, containing 1mM

**Buffer Formulation** EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and

Proclin300.

**Traits** Freeze-dried powder

Purity > 95% Isoelectric Point 8.3

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

## **SEQUENCE**

LEFI AYGTELPSPS YVWFEARFFQ HILHWKPIPN QSESTYYEVA LKQYGNSTWN DIHICRKAQA LSCDLTTFTL DLYHRSYGYR ARVRAVDNSQ YSNWTTTETR FTVDEVILTV DSVTLKAMDG IIYGTIHPPR PTITPAGDEY EQVFKDLRVY KISIRKFSEL KNATKRVKQE TFTLTVPIGV RKFCVKVLPR LESRINKAEW SEEOCLLITT EQYFTVTNLS I

#### **USAGE**


Reconstitute in 100mM NaHCO<sub>3</sub>, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.






Figure. SDS-PAGE

# [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.