

Recombinant Cluster Of Differentiation 73 (CD73) Instruction Manual

SIPB217Bo01

Bos taurus; Bovine (Cattle)

| | |
|---------------------------------|---|
| Source | Prokaryotic expression |
| Host | E.coli |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | Membrane |
| Predicted Molecular Mass | 56.4kDa |
| Accurate Molecular Mass | 56kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Ser48~Val527 with N-terminal His Tag |
| Buffer Formulation | 100mMNaHCO ₃ , 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300. |
| Traits | Freeze-dried powder |
| Purity | > 95% |
| Isoelectric Point | 6.7 |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

SEQUENCE

```
CVNASRCVGG VARLATKVHQ IRRAEHVLL LDAGDQYQGT IWFTVYKGT  
VAHFMNALGY DAMALGNHEF DNGVEGLIDP LLKEVNFPII SANIKAKGPL  
ASKISGLYSP YKILTVGDEV VGIVGYTSKE TPFLSNPGTN LVFEDEITAL  
QPEVDKCLKL NVNKIIALGH SGFEVDKLIQ QKVKGVVQV GGHSTFLYT  
GNPPSKEVPA GQYPFIVTSD DGRKVPVQA YAFGKYLGYL KVEFDEKGNV  
VTSHGNPILL NSSIPEDPNI KADINKWRVK LDNYSTQELG KTIVYLDGTA  
QSCRFRFCNM GNLICDAMIN NNLRHPDEMS WNHVSMCILN GGGIRSPIDE  
RNNGTITWEN LAAVLPFGGT FDLVQLKGST LKKAFEHSVH RYQATGEFL  
QVGGIHVVYD ISRNPGDRVV KLEVLCTQCR VPSYEPLRMD KVKVILPSF  
LVSGGDGFQM IKDEKIKHDS GDQDINV
```

SSK

USAGE

Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.