# Recombinant Clusterin (CLU) Instruction Manual

# SIPB333Hu02

## Homo sapiens (Human)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Membrane, Nucleus, Secreted, Mitochondrion, Cytoplasm,

Endoplasmic reticulum lumen

Predicted Molecular Mass 54.2kDa

Accurate Molecular Mass 54kDa(Analysis of differences refer to the manual)

Residues & Tags 54kDa(Analysis of differences refer to the manual)

Asp23~Arg227 with N-terminal His and GST Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 90% Isoelectric Point 6.6

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

#### **SEQUENCE**

DOTVSDNE LQEMSNQGSK YVNKEIQNAV

NGVKQIKTLI EKTNEERKTL LSNLEEAKKK KEDALNETRE SETKLKELPG VCNETMMALW EECKPCLKQT CMKFYARVCR SGSGLVGRQL EEFLNQSSPF YFWMNGDRID SLLENDRQQT HMLDVMQDHF SRASSIIDEL FQDRFFTREP

QDTYHYLPFS LPHRRPHFFF PKSRIVR

#### USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

#### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



**Image** 

18

22

14

Figure. SDS-PAGE

### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.