# Recombinant Tenascin C (TNC) Instruction Manual

# SIPB376Ra03

# Rattus norvegicus (Rat)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass19.5kDa

Accurate Molecular Mass 25kDa(Analysis of differences refer to the manual)

**Residues & Tags** Val175~Gly342 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 95% Isoelectric Point 4.3

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

# **SEQUENCE**

VCEPGW KGPNCSEPEC PGNCNLRGQC

LDGQCICDQG FTGEDCSQLA CPNDCNDQGK CVNGVCVCFE GYAGLDCGLE VCPVPCSEEH GMCVDGRCVC KDGFAGEDCN EPLCFNNCNN RGRCVENECV

CDEGFTGEDC SELICPNDCF DRGRCINGTC YCEEGFTGED CG

### USAGE

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.




Figure. SDS-PAGE

# [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.