Recombinant Cytokeratin 18 (CK18) Instruction Manual

SIPB410Ra01

Rattus norvegicus (Rat)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Nucleus, Cytoplasm

Predicted Molecular Mass 39.6kDa

Accurate Molecular Mass 40kDa(Analysis of differences refer to the manual)

Residues & Tags Glu72~Gly380 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 5.2

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

EKETMQDLN DRLASYLDKV KNLETENRRL
ESKIREYLEK RGPQGVRDWG HYFKTIEDLR AQIFANSVDN ARIVLQIDNA
RLAADDFRVK YETELAMRQS VESDIHGLRK VVDDTNITRL QLETEIEALK
EELLFMKKNH EEEVQGLEAQ IASSGLTVEV DAPKSQDLSK IMADIRAQYE
QLAQKNREEL DKYWSQQIEE STTVVTTKSA EIRDAETTLL ELRRTLQTLE
IDLDSMKNQN INLENNLGEV EARYRVQMEQ LNGVLLHLES ELAQTRAEGQ
ROTQEYEALL NIKVKLEAEI ATYRRLLEDG

USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.





SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.