

Recombinant Alpha-1-Antitrypsin (a1AT) Instruction Manual

SIPB452Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted, Endoplasmic reticulum lumen
Predicted Molecular Mass	48.0kDa
Accurate Molecular Mass	48kDa(Analysis of differences refer to the manual)
Residues & Tags	Glu25~Lys418 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	5.4
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

EDPQGD AAQKTDTSHH DQDHPTFNKI
TPNLAEFAFS LYRQLAHQSN STNIFFSPVS IATAFAMILSL GTKADTHDEI
LEGLNFnLTe IPEAQIHEGF QELLRTLNQP DSQLQLTTGN GLFLSEGLKL
VDKFLEDVKK LYHSEAFTVN FGDTEEAKKQ INDYVEKGtQ GKIVDLVKEl
DRDTVFALVN YIFFKGKWER PFEVKDTEEE DFHVDQVTTV KVPMMKRLGM
FNIQHCKKLS SWVLLMKYLG NATAIFFLPD EGKLQHLENE LTHDIITKFL
ENEDRRSASL HLPKLSITGT YDLKSVLGQL GITKVFSNGA DLSGVTEEAP
LKLSKAHVKA VLTIDEKGTE AAGAMFLEAI PMSIPPEVKF NKPFVFLMIE
QNTKSPLFMG KVNVNPTQK

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

