Recombinant Mannose Associated Serine Protease 1 (MASP1) Instruction Manual

SIPB575Mu01

Mus musculus (Mouse)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass52.6kDa

Accurate Molecular Mass 53kDa(Analysis of differences refer to the manual)

Residues & Tags His25~Arg453 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01%skl, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 5.1

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

HTVELN EMFGQIQSPG YPDSYPSDSE
VTWNITVPEG FRIKLYFMHF NLESSYLCEY DYVKVETEDQ VLATFCGRET
TDTEQTPGQE VVLSPGTFMS VTFRSDFSNE ERFTGFDAHY MAVDVDECKE
REDEELSCDH YCHNYIGGYY CSCRFGYILH TDNRTCRVEC SGNLFTQRTG
TITSPDYPNP YPKSSECSYT IDLEEGFMVS LQFEDIFDIE DHPEVPCPYD
YIKIKAGSKV WGPFCGEKSP EPISTQTHSV QILFRSDNSG ENRGWRLSYR
AAGNECPKLQ PPVYGKIEPS QAVYSFKDQV LVSCDTGYKV LKDNEVMDTF
QIECLKDGAW SNKIPTCKIV DCGAPAGLKH GLVTFSTRNN LTTYKSEIRY
SCQQPYYKML HNTTGVYTCS AHGTWTNEVL KRSLPTCLPV CGVPKFSRKQ
ISR

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.