# Recombinant Major Histocompatibility Complex Class I E (MHCE) Instruction Manual

# SIPB581Hu01

### Homo sapiens (Human)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass41.5kDa

Accurate Molecular Mass 42kDa(Analysis of differences refer to the manual)

**Residues & Tags** Gly22~Leu358 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 95% Isoelectric Point 5.8

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

### **SEQUENCE**

		<b>GSHSLKYFH</b>	TSVSRPGRGE	PRFISVGYVD
DTQFVRFDND	AASPRMVPRA	<b>PWMEQEGSEY</b>	WDRETRSARD	TAQIFRVNLR
TLRGYYNQSE	AGSHTLQWMH	GCELGPDRRF	LRGYEQFAYD	GKDYLTLNED
LRSWTAVDTA	AQISEQKSND	ASEAEHQRAY	LEDTCVEWLH	KYLEKGKETL
LHLEPPKTHV	THHPISDHEA	TLRCWALGFY	PAEITLTWQQ	DGEGHTQDTE
LVETRPAGDG	TFQKWAAVVV	PSGEEQRYTC	HVQHEGLPEP	VTLRWKPASQ
PTIPIVGIIA	GLVLLGSVVS	GAVVAAVIWR	KKSSGGKGGS	YSKAEWSDSA
OGSESHSI				

### **USAGE**

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.




Figure. SDS-PAGE

## [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.