

Recombinant Thioredoxin- interacting Protein (TXNIP) Instruction Manual

SIPB599Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Cytoplasm
Predicted Molecular Mass	41.1kDa
Accurate Molecular Mass	46kDa(Analysis of differences refer to the manual)
Residues & Tags	Ile32~Ala367 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	7.6
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
                                     IVEVCEVTR VKAVRILACG
VAKVLWMQGS QQCKQTSEYL RYEDTLLLED QPTGENEMVI MRPGNKYEYK
FGFELPQGPL GTSFKGKYGC VDYWKAFD RPSQPTQETK KNFEVVDLVD
VNTPDLMAPV SAKKEKKVSC MFIPDGRVSV SARIDRKGFC EGDEISIHAD
FENTCSRIVV PKAAIVARHT YLANGQTKVL TQKLSSVRGN HIISGTCASW
RGKSLRVQKI RPSILGCNII RVEYSLLIYV SVPGSKKVIL DLPLVIGSRS
GLSSRTSSMA SRTSSEMSW DLNIPDTPEA PPCYMDVIPE DHRLESPTTP
LLDDMDGSQD SPIFMYA
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.