Recombinant VGF Nerve Growth Factor Inducible (VGF) Instruction Manual

SIPB602Mu02

Mus musculus (Mouse)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1μg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass36.5kDa

Accurate Molecular Mass 43kDa(Analysis of differences refer to the manual)

Residues & Tags Arg284~Arg576 with N-terminal His Tag

Buffer Formulation PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5%

Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 95% Isoelectric Point 4.2

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

RLEGSFL GGSEAGERLL QQGLAQVEAG RRQAEATRQA AAQEERLADL ASDLLLQYLL QGGARQRDLG GRELQETQQE RENEREEEAE QERRGGGEDD VGEEDEEAAE AEAEAEEAER ARQNALLFAE EEDGEAGAED KRSQEEAPGH RRKDAEGAEE GGEEDDDDEE MDPQTIDSLI ELSTKLHLPA DDVVSIIEEV EEKRKRKNA PPEPVPPPRA APAPTHVRSP QPPPPAPARD ELPDWNEVLP PWDREEDEVF PPGPYHPFPN

YIRPRTLQPP ASSRRRHFHH ALPPAR

USAGE


Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.