

Recombinant Hyaluronidase (HAase) Instruction Manual

SIPB610Hu02

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1μg (determined by the LAL method)
Subcellular Location	Secreted, Lysosome
Predicted Molecular Mass	49.7kDa
Accurate Molecular Mass	46kDa(Analysis of differences refer to the manual)
Residues & Tags	Phe22~Trp435 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	6.9
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

FRGPLLPNR PFTTVWNANT QWCLERHGVD
VDVSVF DVVA NPGQTFRGPD MTIFYSSQLG TYPYYTPTGE PVFGGLPQNA
SLIAHLARTF QDILAAIPAP DFSGLAVIDW EAWRPRWAFN WDTKDIYRQR
SRALVQAQHP DWPA PAPQVEAV AQDQFQGAAR AWMAGTLQLG RALRPRGLWG
FYGF PD CNY DFLSP NYTGQ CPSGIR AQND QLGWLWGQSR ALYPSIYMPA
VLEG TGKSQM YVQHRV AEAF RVAVAAGDPN LPVLPYVQIF YDTTNHFLPL
DELEHSLGES AAQGAAGVVL WWSWENTRTK ESCQAIKEYM DTTLGP FILN
VTSGALLCSQ ALCSGHGR CV RRTSHPK ALL LLNPASFSIQ LTPGGGPLSL
RGALSLEDQA QMAVEFKCRC YPGWQAPWCE RKSMW

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

