Recombinant Cluster Of Differentiation 164 (CD164) Instruction Manual

SIPB674Mu01

Mus musculus (Mouse)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass18.5kDa

Accurate Molecular Mass 19kDa(Analysis of differences refer to the manual)

Residues & Tags Gln24~Asp162 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 6.7

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

QPNITTL APNVTEVPTT TTKVVPTTQM
PTVLPETCAS FNSCVSCVNA TFTNNITCFW LHCQEANKTY CANEPLSNCS
QVNRTDLCSV IPPTTPVPTN STAKPTTRPS SPTPTPSVVT SAGTTNTTLT
PTSQPERKST FD

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



15% SDS-PAGE SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.