# Recombinant Gamma Actin (ACTG) Instruction Manual

# SIPB759Mu01

## Mus musculus (Mouse)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationCytoplasmPredicted Molecular Mass43.1kDa

Accurate Molecular Mass 43kDa(Analysis of differences refer to the manual)

Residues & Tags Met1~Phe375 with N-terminal His Tag

**Buffer Formulation** PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

**Traits** Freeze-dried powder

Purity > 95% Isoelectric Point 5.2

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

#### **SEQUENCE**

MEEEIAALVI DNGSGMCKAG FAGDDAPRAV FPSIVGRPRH QGVMVGMGQK DSYVGDEAQS KRGILTLKYP IEHGIVTNWD DMEKIWHHTF YNELRVAPEE HPVLLTEAPL NPKANREKMT QIMFETFNTP AMYVAIQAVL SLYASGRTTG IVMDSGDGVT HTVPIYEGYA LPHAILRLDL AGRDLTDYLM KILTERGYSF TTTAEREIVR DIKEKLCYVA LDFEQEMATA ASSSSLEKSY ELPDGQVITI GNERFRCPEA LFQPSFLGME SCGIHETTFN SIMKCDVDIR KDLYANTVLS GGTTMYPGIA DRMQKEITAL APSTMKIKII APPERKYSVW IGGSILASLS

TFQQMWISKQ EYDESGPSIV HRKCF

#### USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

#### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



(A TOTAL ASSOCIATION OF PURSUITATION OF A REPORT OF A TOTAL ASSOCIATION OF A TOTAL ASSOCIAT




Figure. SDS-PAGE

### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.