Recombinant Adenylate Cyclase 2, Brain (ADCY2) Instruction Manual

SIPB764Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass29.6kDa

Accurate Molecular Mass 31kDa(Analysis of differences refer to the manual)

Residues & Tags Ala294~Thr525 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 8.6

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

				ADIVGFT
RLASDCSPGE	LVHMLNELFG	KFDQIAKENE	CMRIKILGDC	YYCVSGLPIS
LPNHAKNOVK	MGLDMCEAIK	KVRDATGVDI	NMRVGVHSGN	VLCGVIGLQK
WQYDVWSHDV	TLANHMEAGG	VPGRVHISSV	TLEHLNGAYK	VEEGDGDIRD
PYLKQHLVKT	YFVINPKGER	RSPQHLFRPR	HTLDGAKMRA	SVRMTRYLES
WGAAKPFAHL	HHRDSMTTEN	GKIST		

USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.