Recombinant Enolase 1 (ENO1) Instruction Manual

SIPB782Bo01

Bos taurus; Bovine (Cattle)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Membrane, Cytoplasm

Predicted Molecular Mass 48.9kDa

Accurate Molecular Mass 48kDa(Analysis of differences refer to the manual)

Residues & Tags Met1~Lys434 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 6.5

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

MSILKVHARE IFDSRGNPTV EVDLFTAKGL FRAAVPSGAS TGIYEALELR
DNDKTRYMGK GVSKAVEHIN KTIAPALVSK KLNVVEQEKI DKLMIEMDGT
ENKSKFGANA ILGVSLAVCK AGAVEKGVPL YRHIADLAGN AEVILPVPAF
NVINGGSHAG NKLAMQEFMI LPVGAENFRE AMRIGAEVYH NLKNVIKEKY
GKDATNVGDE GGFAPNILEN KEALELLKNA IGKAGYSDKV VIGMDVAASE
FYRSGKYDLD FKSPDDPSRY ITPDELANLY KSFIRDYPVV SIEDPFDQDD
WEAWQKFTAS AGIQVVGDDL TVTNPKRIAK AVSEKSCNCL LLKVNQIGSV
TESLQACKLA QSNGWGVMVS HRSGETEDTF IADLVVGLCT GQIKTVAPCR
SERLAKYNQI LRIEEELGSK AKFAGRSFRN PLAK

USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

SDS-PAGE Image

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.