Recombinant Interleukin 2 Receptor Gamma (IL2Rg) Instruction Manual

SIPB802Mu01

Mus musculus (Mouse)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass29.5kDa

Accurate Molecular Mass 29kDa(Analysis of differences refer to the manual)

Residues & Tags Trp23~Ala263 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 95% Isoelectric Point 7.2

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

WSSKVLMS SANEDIKADL ILTSTAPEHL

SAPTLPLPEV QCFVFNIEYM NCTWNSSSEP QATNLTLHYR YKVSDNNTFQ ECSHYLFSKE ITSGCQIQKE DIQLYQTFVV QLQDPQKPQR RAVQKLNLQN LVIPRAPENL TLSNLSESQL ELRWKSRHIK ERCLQYLVQY RSNRDRSWTE LIVNHEPRFS LPSVDELKRY TFRVRSRYNP ICGSSQQWSK WSQPVHWGSH

TVEENPSLFA LEA

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE


Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image

GATTERETTIANTITIAN TERRETTANDE STITETTIAN TITETTIAN TITETTIAN TITETTIAN TERRETTAN TERRETTANDE STANDARDE ST



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.