Recombinant Integrin Beta 2 (CD18) Instruction Manual

SIPB817Hu03

Homo sapiens (Human)

Prokaryotic expression
E.coli
<1.0EU per 1µg (determined by the LAL method)
Membrane
42.0kDa
44kDa(Analysis of differences refer to the manual)
Gln23~Val367 with N-terminal His Tag
PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.
Freeze-dried powder
> 90%
6.6
Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

QECTKFKV SSCRECIESG PGCTWCQKLN FTGPGDPDSI RCDTRPQLLM RGCAADDIMD PTSLAETQED HNGGQKQLSP QKVTLYLRPG QAAAFNVTFR RAKGYPIDLY YLMDLSYSML DDLRNVKKLG GDLLRALNEI TESGRIGFGS FVDKTVLPFV NTHPDKLRNP CPNKEKECQP PFAFRHVLKL TNNSNQFQTE VGKQLISGNL DAPEGGLDAM MQVAACPEEI GWRNVTRLLV FATDDGFHFA GDGKLGAILT PNDGRCHLED NLYKRSNEFD YPSVGQLAHK LAENNIQPIF AVTSRMVKTY EKLTEIIPKS AVGELSEDSS NVVQLIKNAY NKLSSRV

USAGE

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

Image