# **Recombinant Interferon Regulatory Factor 3 (IRF3) Instruction Manual**

# SIPB819Hu01

### Homo sapiens (Human)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Nucleus, Cytoplasm

Predicted Molecular Mass 43.8kDa

Accurate Molecular Mass 44kDa(Analysis of differences refer to the manual)

Residues & Tags Met1~Lys360 with N-terminal His Tag

20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL,

5% Trehalose.

**Traits** Freeze-dried powder

Purity > 95% Isoelectric Point 5.8

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

## **SEQUENCE**

**Buffer Formulation** 

MGTPKPRILP WLVSQLDLGQ LEGVAWVNKS RTRFRIPWKH GLRQDAQQED FGIFQAWAEA TGAYVPGRDK PDLPTWKRNF RSALNRKEGL RLAEDRSKDP HDPHKIYEFV NSGVGDFSQP DTSPDTNGGG STSDTQEDIL DELLGNMVLA PLPDPGPPSL AVAPEPCPQP LRSPSLDNPT PFPNLGPSEN PLKRLLVPGE EWEFEVTAFY RGRQVFQQTI SCPEGLRLVG SEVGDRTLPG WPVTLPDPGM SLTDRGVMSY VRHVLSCLGG GLALWRAGQW LWAQRLGHCH TYWAVSEELL PNSGHGPDGE VPKDKEGGVF DLGPFIVDLI TFTEGSGRSP RYALWFCVGE SWPODOPWTK

#### USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

#### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



# [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.