# Recombinant Lymphocyte Function Associated Antigen 1 Alpha (CD11a) Instruction Manual

## SIPB971Hu01

#### Homo sapiens (Human)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass32.5kDa

Accurate Molecular Mass 33kDa(Analysis of differences refer to the manual)

**Residues & Tags** Asn809~His1066 with N-terminal His Tag

Buffer Formulation PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5%

Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 95% Isoelectric Point 5.5

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

#### **SEQUENCE**

NL EEDAYWVQLD LHFPPGLSFR KVEMLKPHSQ IPVSCEELPE ESRLLSRALS CNVSSPIFKA GHSVALQMMF NTLVNSSWGD SVELHANVTC NNEDSDLLED NSATTIIPIL YPINILIQDQ EDSTLYVSFT PKGPKIHQVK HMYQVRIQPS IHDHNIPTLE AVVGVPQPPS EGPITHQWSV QMEPPVPCHY EDLERLPDAA EPCLPGALFR CPVVFRQEIL VQVIGTLELV GEIEASSMFS LCSSLSISFN SSKHFH

#### USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

#### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



TEXT TO CONSISTENCE THAT THE THAT THE SECOND TO THE THE THE SECOND SECON




Figure. SDS-PAGE

### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.