

Recombinant Transferrin (TF) Instruction Manual

SIPC020Ra01

Rattus norvegicus (Rat)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	42.3kDa
Accurate Molecular Mass	45kDa(Analysis of differences refer to the manual)
Residues & Tags	Val360~Lys683 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	6.6
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
V KWCALSHQER AKCDEWSVSS NGQIECESAE STEDCIDKIV  
NGEADAMSLD GGHAYIAGQC GLVPVMAENY DISSCTNPQS DVFPKGYAV  
AVVKASDSSI NWNMLKGGKS CHTGVDRTAG WNIPMGLLFS RINHCKFDEF  
FSQGCAPGYK KNSTLCDLCI GPAKCAPNMR EGYNGYTGAF QCLVEKGDVA  
FVKHQTVLEN TNGKNTAAWA KDLKQEDFQL LCPDGTKKPV TEFATCHLAQ  
APNHVVVSRK EKAARVSTVL TAQKDLFWKG DKDCTGNFCL FRSSTKDLLF  
RDDTKCLTKL PEGTTYEAYL GAEYLQAVGN IRK
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.