# Recombinant Laminin Gamma 2 (LAMC2) Instruction Manual

## SIPC054Hu03

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	49.6kDa
Accurate Molecular Mass	50kDa(Analysis of differences refer to the manual)
Residues & Tags	Cys416~Cys602 with N-terminal His and GST Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	5.5
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

### SEQUENCE

CNCQG GGACDPDTGD CYSGDENPDI ECADCPIGFY NDPHDPRSCK PCPCHNGFSC SVMPETEEVV CNNCPPGVTG ARCELCADGY FGDPFGEHGP VRPCQPCQCN NNVDPSASGN CDRLTGRCLK CIHNTAGIYC DQCKAGYFGD PLAPNPADKC RACNCNPMGS EPVGCRSDGT CVCKPGFGGP NC

### USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


### STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

#### STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### Image



### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.