

Recombinant Integrin Beta 6 (ITGb6) Instruction Manual

SIPC066Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Membrane
Predicted Molecular Mass	30.5kDa
Accurate Molecular Mass	29kDa(Analysis of differences refer to the manual)
Residues & Tags	Tyr131~Leu371 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	4.8
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
TIKELGSRLS KEMSKLTSNF RLGFGSFVEK YPVDLYYLM DLSASMDDL N  
PYFCLPTFGF KHILPLTND AERFNEIVKN QKISANIDTP EGGFDAIMQA A  
VCKEKIGWRN DSLHLLVFS DADSHFGMDS KLAGIVIPND GLCHLDSKNE  
YSMSTVLEYP TIGQLIDKLV QNNVLLIFAV TQEQVHLYEN YAKLIPGATV  
GLLQKDSGNI LQLIISAYEE L
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE


Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image

TCCTGGGTTTGATTTCCTATGGACTCTGGCTCGGATCCGCTTAAAGCAATGAGATGGGCTGGGCTGGGTTTCAGGAGTGTCTAAATAGCGAATTTCAGCTGGGTTTGGACTTTGGGCTTTTGTGCAAAAGCTGGTTCCTTCTTTCGCAAGCAAGTTGCAGCCCTTGCACTAT
Y P V D I Y V L W D I S A S M D D D L W T I K E L G S R I S K E H S E L T S W F R I G F G S F V E K P V S P F V K T T P E E I A N P C S S I



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.