

Recombinant Flotillin 2 (FLOT2) Instruction Manual

SIPC132Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Membrane
Predicted Molecular Mass	50.7kDa
Accurate Molecular Mass	55kDa(Analysis of differences refer to the manual)
Residues & Tags	Met1~Val428 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	4.9
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

MGNCHTVGPN EALVVSGGCC GSDYKQYVFG GWAIAWWCIS DTQRISLEIM
TLQPRCEDVE TAEGVALTVT GVAQVKIMTE KELLAVACEQ FLGKNVQDIK
NVVLQTLEGH LRSILGTLTV EQIYQDRDQF AKLVREVAAP DVGRMGIEIL
SFTIKDVYDK VDYLSSLGKT QTAVVQRDAD IGVAEAERDA GIRAECKKE
MLDVKFMADT KIADSKRAFE LQKSAFSEEV NIKTAEAQLA YELQGAREQQ
KIRQEEIEIE VVQRKKQIAV EAQEILRTDK ELIATVRRPA EAEAHRIQQI
AEGEKVKQVL LAQAEAEKIR KIGEAEAAVI EAMGKAEAER MKLKAЕAYQK
YGDAAKMALV LEALPQIAAK IAAPLTKVDE IVVLSGDNISK VTSEVNRLLA
ELPASVHALT GVDLSKIPLI KKATGVQV

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

