Recombinant Furin (FUR) Instruction Manual

SIPC359Mu02

Mus musculus (Mouse)

| Source | Prokaryotic expression |
|---------------------------|---|
| Host | E.coli |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | Chromosome |
| Predicted Molecular Mass | 49.2kDa |
| Accurate Molecular Mass | 52kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Leu578~Phe734 with N-terminal His and GST Tag |
| Buffer Formulation | 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300. |
| Traits | Freeze-dried powder |
| Purity | > 97% |
| Isoelectric Point | 6.0 |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

SEQUENCE

LST PPESSGCKTL TSSQACVVCE EGYSLHQKSC VQHCPPGFIP QVLDTHYSTE NDVEIIRASV CTPCHASCAT CQGPAPTDCL SCPSHASLDP VEQTCSRQSQ SSRESRPQQQ PPALRPEVEM EPRLQAGLAS HLPEVLAGLS CLIIVLIFGI VFLF

USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.