Recombinant Calpain 10 (CAPN10) Instruction Manual

SIPC368Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted
Predicted Molecular Mass	40.2kDa
Accurate Molecular Mass	40kDa(Analysis of differences refer to the manual)
Residues & Tags	Thr322~Gln654 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	9.6
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

TEAGHLQSL YTERLLCHTR ALPGAWVKGQ SAGGCRNNSG FPSNPKFWLR VSEPSEVYIA VLQRSRLHAA DWAGRARALV GDSHTSWSPA SIPGKHYQAV GLHLWKVEKR RVNLPRVLSM PPVAGTACHA YDREVHLRCE LSPGYYLAVP STFLKDAPGE FLLRVFSTGR VSLSAIRAVA KNTTPGAALP AGEWGTVQLR GSWRVGQTAG GSRNFASYPT NPCFPFSVPE GPGPRCVRIT LHQHCRPSDT EFHPIGFHIF QVPEGGRSQD APPLLLQEPL LSCVPHRYAQ EVSRLCLLPA GTYKVVPSTY LPDTEGAFTV TIATRIDRPS IHSQ

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

STABILITY

Image

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

kDa 70 44 33 26 22 18 14 10

Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.