

Recombinant Tumor Necrosis Factor Alpha Induced Protein 3 (TNF α IP3) Instruction Manual

SIPC463Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1 μ g (determined by the LAL method)
Subcellular Location	Nucleus, Cytoplasm
Predicted Molecular Mass	38.9kDa
Accurate Molecular Mass	39kDa(Analysis of differences refer to the manual)
Residues & Tags	Met1~Glu300 with N-terminal His Tag
Buffer Formulation	100mMNaHCO ₃ , 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	8.6
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
MAEQVLPQAL YLSNMRKAVK IRERTPEDIF KPTNGIIHHF KTMHRYTLEM  
FRTCQFCPQF REIIHKALID RNIQATLESQ KKLNWCREVR KLVALKTNQD  
GNCLMHATSQ YMWGVQDIDL VLRKALFSTL KETDTRNFKF RWQLESLSQ  
EFVETGLCYD TRNWNDEWDN LIKMASTDTP MARSGLQYNS LEEIHIFVLC  
NILRRPIIVI SDKMLRSLES GSNFAPLKVG GIYLPLHWA QECYRYPIVL  
GYDSHHFVPL VTLKDSGPEI RAVPLVNRDR GRFEDLKVHF LDPENEMKE
```

USAGE

Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.