

Recombinant Sortilin 1 (SORT1) Instruction Manual

SIPC575Hu01

Homo sapiens (Human)

| | |
|---------------------------------|---|
| Source | Prokaryotic expression |
| Host | E.coli |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | Membrane |
| Predicted Molecular Mass | 59.7kDa |
| Accurate Molecular Mass | 60kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Trp50~Asp320 with N-terminal His and GST Tag |
| Buffer Formulation | 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose. |
| Traits | Freeze-dried powder |
| Purity | > 90% |
| Isoelectric Point | 8.7 |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

SEQUENCE

SGPIGVSWGL RAAAAGGAFP RGGRWRRSAP GEDEECGRVR DFVAKLANNT
HQHFVDDLRLG SVSLSWVGDS TGVILVLTTF HVPLVIMTFG QSKLYRSEDY
GKNFKDITDL INNTFIRTEF GMAIGPENSG KVVLTAEVSG GSRGGRIFRS
SDFAKNFVQT DLPFHPLTQM MYSPQNSDYL LALSTENGLW VSKNFGGKWE
EIHKAVCLAK WGSDNITIFFT TYANGSCKAD LGALELWRTS DLGKSFKTIG
VKIYSFGLGG RFLFASVMAD

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.