# Recombinant Histidyl tRNA Synthetase (HARS) Instruction Manual

# SIPC577Hu01

### Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Cytoplasm
Predicted Molecular Mass	37.3kDa
Accurate Molecular Mass	37kDa(Analysis of differences refer to the manual)
Residues & Tags	Met1~Gln294 with N-terminal His Tag
Buffer Formulation	100mMNaHCO <sub>3</sub> , 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	8.1
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

#### SEQUENCE

MAERAALEEL VKLQGERVRG LKQQKASAEL IEEEVAKLLK LKAQLGPDES KQKFVLKTPK GTRDYSPRQM AVREKVFDVI IRCFKRHGAE VIDTPVFELK ETLMGKYGED SKLIYDLKDQ GGELLSLRYD LTVPFARYLA MNKLTNIKRY HIAKVYRRDN PAMTRGRYRE FYQCDFDIAG NFDPMIPDAE CLKIMCEILS SLQIGDFLVK VNDRRILDGM FAICGVSDSK FRTICSSVDK LDKVSWEEVK NEMVGEKGLA PEVADRIGDY VQQHGGVSLV EQLLQDPKLS QNKQ

#### USAGE

Reconstitute in 100mM NaHCO3, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


#### STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

#### STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### Image



SDS-PAGE Image

## [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.