# **Recombinant Integrin Alpha 11** (ITGa11) Instruction Manual

# SIPC606Hu01

#### Homo sapiens (Human)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass38.1kDa

Accurate Molecular Mass 38kDa(Analysis of differences refer to the manual)

**Residues & Tags** Phe23~His328 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 95% Isoelectric Point 8.4

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

## **SEQUENCE**

		FNMDTRKP	RVIPGSRTAF	FGYTVQQHDI
SGNKWLVVGA	PLETNGYQKT	GDVYKCPVIH	GNCTKLNLGR	VTLSNVSERK
DNMRLGLSLA	TNPKDNSFLA	CSPLWSHECG	SSYYTTGMCS	RVNSNFRFSK
TVAPALQRCQ	TYMDIVIVLD	GSNSIYPWVE	VQHFLINILK	KFYIGPGQIQ
VGVVQYGEDV	VHEFHLNDYR	SVKDVVEAAS	HIEQRGGTET	RTAFGIEFAR
SEAFQKGGRK	GAKKVMIVIT	DGESHDSPDL	EKVIQQSERD	NVTRYAVAVL
GYVNRRGINE	FTFI NFTKYT	ASDPODKH		

## USAGE


Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

#### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



## Figure. SDS-PAGE

#### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.