

Recombinant Neutrophil Cytosolic Factor 1 (NCF1) Instruction Manual

SIPC622Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Membrane, Cytoplasm
Predicted Molecular Mass	48.4kDa
Accurate Molecular Mass	48kDa(Analysis of differences refer to the manual)
Residues & Tags	Met1~Val390 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	9.2
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
MGDTFIRHIA LLGFEKRFVP SQHYVYMLV KWQDLSEKVV YRRFTEIYEF
HKTLKEMFPI EAGAINPENR IIPHLPAKWK FDGQRAAENR QGTLTEYCGT
LMSLPTKISR CPHLLDFFKV RPDDLKLPD NQTKKPETYL MPKDGKSTAT
DITGPIILQT YRAIANYEKT SGSEMALSTG DVVEVVEKSE SGWWFCQMK
KRGWIPASFL EPLDSPDETE DPEPNYAGEP YVAIKAYTAV EGDEVSLL
EAVEVIHKLL DGWVWIRKDD VTGYFPSMYL QKSGQDVSQA QRQIKRGAPP
RRSSIRNAHS IHQRSRKRLS QDAYRRNSVR FLQRRRRQAR PGPQSPGSPL
EEERQTQRSK PQPAVPPRPS ADLILNRCSE STKRKLASAV
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



15% SDS-PAGE
SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.