Recombinant Embryonic Ectoderm Development (EED) Instruction Manual

SIPC637Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Nucleus, Cytoplasm

Predicted Molecular Mass 38.4kDa

Accurate Molecular Mass 38kDa(Analysis of differences refer to the manual)

Residues & Tags Asp142~Arg441 with N-terminal His Tag

100mMNaHCO₃, 500mMNaCl, pH8.3, containing 1mM

Buffer Formulation EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and

Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 6.9

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

				DADENFYTC
AWTYDSNTSH	PLLAVAGSRG	IIRIINPITM	QCIKHYVGHG	NAINELKFHP
RDPNLLLSVS	KDHALRLWNI	QTDTLVAIFG	GVEGHRDEVL	SADYDLLGEK
IMSCGMDHSL	KLWRINSKRM	MNAIKESYDY	NPNKTNRPFI	SQKIHFPDFS
TRDIHRNYVD	CVRWLGDLIL	SKSCENAIVC	WKPGKMEDDI	DKIKPSESNV
TILGRFDYSQ	CDIWYMRFSM	DFWQKMLALG	NQVGKLYVWD	LEVEDPHKAK
CTTLTHHKCG	AAIRQTSFSR	DSSILIAVCD	DASIWRWDRL	R

USAGE

Reconstitute in 100mM NaHCO3, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.