

Recombinant Nicastrin (NCSTN) Instruction Manual

SIPC651Ra01

Rattus norvegicus (Rat)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Membrane
Predicted Molecular Mass	59.1KDa
Accurate Molecular Mass	59kDa(Analysis of differences refer to the manual)
Residues & Tags	Val35~Gln534 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	5.2
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

VERKIY IPLNKTAPCV
RLLNATHQIG CQSSISGDTG VIHVVEKEDD LKWVLTGPN PPYMVLLLEGK
LFTRDIMEKL KGETSRIAGL AVTLAKPNST SSFSPSVQCP NDGFGIYSNS
YGPEFAHCKK TLWNELGNGL AYDDFSFPIF LLEDENETKV IKQCYQDHNL
GQNGSAPSFP LCAMQLFSHM HAVISTATCM RRSFIQSTFS INPEIVCDPL
SDYNVWSMLK PINTSGGLEP DVRVVAATR LDSRSFFWNV APGAESAVAS
FVTQLAAAEA LHKAPDVTTL PRNVMFVFFQ GETFDYIGSS RMVYDMENK
FPVRLNIDS FVELGQVALR TSLELWMHTD PMSQKNESVK NQVEDLLVTL
EQSGADTPQV VLSRLVQSQA LPPSSLQRFL RARNISGVVL ADHSGSFHNR
YYQSIYDTAE NINVTYPESQ SPEEDLNFTV DTAKALADVA TVLARALYKL
AGGTNFNNSI QADPQTVTRL LYGFLVRANN SWFQ

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.
Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

