# Recombinant Thyroid Stimulating Hormone Receptor (TSHR) Instruction Manual

# SIPC676Hu01

### Homo sapiens (Human)

**Source** Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass48.3kDa

Accurate Molecular Mass 50kDa(Analysis of differences refer to the manual)

**Residues & Tags** Gly21~Gly413 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 95% Isoelectric Point 5.4

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

### **SEQUENCE**

GMGCSSPPCE CHQEEDFRVT CKDIQRIPSL
PPSTQTLKLI ETHLRTIPSH AFSNLPNISR IYVSIDVTLQ QLESHSFYNL
SKVTHIEIRN TRNLTYIDPD ALKELPLLKF LGIFNTGLKM FPDLTKVYST
DIFFILEITD NPYMTSIPVN AFQGLCNETL TLKLYNNGFT SVQGYAFNGT
KLDAVYLNKN KYLTVIDKDA FGGVYSGPSL LDVSQTSVTA LPSKGLEHLK
ELIARNTWTL KKLPLSLSFL HLTRADLSYP SHCCAFKNQK KIRGILESLM
CNESSMQSLR QRKSVNALNS PLHQEYEENL GDSIVGYKEK SKFQDTHNNA
HYYVFFEEQE DEIIGFGQEL KNPQEETLQA FDSHYDYTIC GDSEDMVCTP
KSDEFNPCED IMG

### USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.





### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.