

Recombinant Cytochrome P450 17A1 (CYP17A1) Instruction Manual

SIPC686Mu01

Mus musculus (Mouse)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Membrane
Predicted Molecular Mass	38.6kDa
Accurate Molecular Mass	38kDa(Analysis of differences refer to the manual)
Residues & Tags	Thr201~Thr507 with N-terminal His Tag
Buffer Formulation	100mMNaHCO ₃ , 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	5.6
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
TEGIVDVLGH SDLVDIFPWL KIFPNKNLEM IKEHTKIREK TLVEMFEKCK  
EKFNSESLSS LTDILIQAKM NAENNNTGEG QDPSVFSDKH ILVTVGDI  
AGIETTSSVL NWILAFVHN PEVKRKIQKE IDQYVGFST PSFNDRTHLL  
MLEATIREVL RIRPVAPLLI PHKANIDSSI GEFAIPKDTL VIINLWALHH  
DKNEWDQPDR FMPERFLDPT GSHLITPTPS YLPGAGPRS CIGEARARQE  
LFIFMALLLQ RFDVSDDK QLPCLVGDPK VVFLIDPFKV KITVRQAWKD  
AQVEVST
```

USAGE

Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

