Recombinant Purine Nucleoside Phosphorylase (PNP) Instruction Manual

SIPC853Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationCytoplasmPredicted Molecular Mass35.8kDa

Accurate Molecular Mass 36kDa(Analysis of differences refer to the manual)

Residues & Tags Met1~Ser289 with N-terminal His Tag

100mMNaHCO₃, 500mMNaCl, pH8.3, containing 1mM

Buffer Formulation EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and

Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 6.5

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

MENGYTYEDY KNTAEWLLSH TKHRPQVAII CGSGLGGLTD KLTQAQIFDY GEIPNFPRST VPGHAGRLVF GFLNGRACVM MQGRFHMYEG YPLWKVTFPV RVFHLLGVDT LVVTNAAGGL NPKFEVGDIM LIRDHINLPG FSGQNPLRGP NDERFGDRFP AMSDAYDRTM RQRALSTWKQ MGEQRELQEG TYVMVAGPSF ETVAECRVLQ KLGADAVGMS TVPEVIVARH CGLRVFGFSL ITNKVIMDYE SLEKANHEEV LAAGKQAAQK LEQFVSILMA SIPLPDKAS

USAGE

Reconstitute in 100mM NaHCO3, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.