Recombinant Tec Protein Tyrosine Kinase (TEC) Instruction Manual

SIPC888Hu01

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular Location Membrane, Cytoplasm

Predicted Molecular Mass 33.3kDa

Accurate Molecular Mass 33kDa(Analysis of differences refer to the manual)

Residues & Tags Leu370~Val623 with N-terminal His Tag

100mMNaHCO₃, 500mMNaCl, pH8.3, containing 1mM

Buffer Formulation EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and

Proclin300.

Traits Freeze-dried powder

Purity > 97% Isoelectric Point 7.7

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

L TFMRELGSGL FGVVRLGKWR AQYKVAIKAI
REGAMCEEDF IEEAKVMMKL THPKLVQLYG VCTQQKPIYI VTEFMERGCL
LNFLRQRQGH FSRDVLLSMC QDVCEGMEYL ERNSFIHRDL AARNCLVSEA
GVVKVSDFGM ARYVLDDQYT SSSGAKFPVK WCPPEVFNYS RFSSKSDVWS
FGVLMWEVFT EGRMPFEKYT NYEVVTMVTR GHRLYQPKLA SNYVYEVMLR
CWQEKPEGRP SFEDLLRTID ELV

CHQLIN Edin SI EDEENIED EE

USAGE

Reconstitute in ddH₂O to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



15% SDS-PAGE SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.