

# **Recombinant Cholinergic Receptor, Nicotinic, Beta 4 (CHRN<sub>b</sub>4) Instruction Manual**

**SIPD089Hu01**

**Homo sapiens (Human)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Membrane
<b>Predicted Molecular Mass</b>	28.8kDa
<b>Accurate Molecular Mass</b>	32kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Arg22~Thr236 with N-terminal His Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 90%
<b>Isoelectric Point</b>	6.8
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## **SEQUENCE**

RVANAEKKL MDDLLNKTRY NNLIRPATSS  
SQLISIKLQL SLAQLISVNE REQIMTTNVW LKQEWTDYRL TWNSSRYEGV  
NILRIPAKRI WLPDIVLYNN ADGTYEVSVY TNLIVRSNGS VLWLPPAIYK  
SACKIEVKYF PFDQQNCTLK FRSWTDHTE IDMVLMTPTA SMDDFTPSGE  
WDIVALPGRR TVNPQDPSYV DVTYDFIIKR KPLFYT

## **USAGE**

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

## **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### Image

G A T T C G G A T C C G C G A C C C A T T T G C T G T C C A C C A G T C A T G C T A G C C A T A T G G C T G C C G C G C G G C A C C A G G C C G C T G C T G T G A T G




Figure. SDS-PAGE

### **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.